## CORRECTION

**Open Access** 

**Cancer Cell International** 



## Correction to: SIRT6/HIF-1a axis promotes papillary thyroid cancer progression by inducing epithelial–mesenchymal transition

Zhou Yang<sup>1†</sup>, Weiping Yu<sup>1†</sup>, Renhong Huang<sup>1</sup>, Min Ye<sup>1\*</sup> and Zhijun Min<sup>1\*</sup>

Correction to: *Cancer Cell Int* (2019)19:17 https://doi.org/10.1186/s12935-019-0730-4

In the article [1], the authors have found an error of  $\alpha$ -tubulin in Fig. 4d. This error was caused by the same label name, and we put a same picture of  $\alpha$ -tubulin in both BCPAP-NC and BCPAP-SIRT6 group in Fig. 4d during figure processing. The error doesn't affect any results and conclusions of this research. The correct Fig. 4d is given in this correction.

<sup>†</sup>Zhou Yang and Weiping Yu contributed equally to this work.

The online version of the original article can be found at https://doi. org/10.1186/s12935-019-0730-4.

\*Correspondence: Min Ye yemincool@126.com Zhijun Min minzhijun@126.com <sup>1</sup>Department of General Surgery, Shanghai Pudong Hospital, Fudan University Pudong Medical Center, 2800 Gongwei Road, Huinan Town, 201399 Pudong, Shanghai, China



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Dublic Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



Fig. 4 a mRNA expression of HIF-1 $\alpha$  fold to negative control under normoxia and hypoxia. The comparative cycle threshold values (2– $\Delta\Delta$ Ct) and t-test were adopted to analyze the final results. **b** Protein expression of HIF-1 $\alpha$  in each group under normoxia and hypoxia. **c** HIF-1 $\alpha$  expression in TPC1-SIRT6 and TPC1-NC cells after withdrawing CoCl2 for 0–90 min. **d** HIF-1 $\alpha$  expression in BCPAP-SIRT6 and BCPAP-NC cells after withdrawing CoCl2 for 0–90 min. (\*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001)

Accepted: 1 August 2023 Published online: 22 August 2023

## References

 Yang Z, Yu W, Huang R, Ye M, Min Z. SIRT6/HIF-10 axis promotes papillary thyroid cancer progression by inducing epithelial-mesenchymal transition. Cancer Cell Int. 2019;19:17.

## **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.